

pp.). Observed basking on a log in a wetland surrounded mostly by mixed hardwood forest at 1723 h.

Submitted by **JUSTIN T. OGUNI**, College of Veterinary Medicine, University of Georgia, Athens, Georgia 30602, USA; e-mail: goonie16@gmail.com.

SQUAMATA — LIZARDS

HEMIDACTYLUS TURCICUS (Mediterranean Gecko). USA: ALABAMA: TUSCALOOSA Co.: Northport, 3550 Watermelon Road (33.238706°N, 87.552747°W; WGS84). 25 September 2009. Christopher Thawley. University of Alabama Herpetological Collection (UAHC 16363). Verified by Kenneth L. Krysko. New county record (Mount 1975. *Amphibians and Reptiles of Alabama*. Auburn University Agricultural Experiment Station, Auburn, Alabama. 347 pp.). The nearest known record is from Jefferson Co., Alabama (Marion and Bosworth 1982. *Herpetol. Rev.* 13:52) 74 km NE. Juvenile, captured by hand, observed along with two adults, suggesting presence of potential breeding population. Anecdotal reports place additional populations in some dormitories on the University of Alabama main campus in Tuscaloosa, less than 3 km S and in Cottondale, ca. 10 km SE.

Submitted by **CHRISTOPHER J. THAWLEY**, Department of Biological Sciences, University of Alabama, Box 870345, Tuscaloosa, Alabama 35487, USA; e-mail: cjthawley@crimson.ua.edu.

LEIOCEPHALUS CARINATUS (Northern Curly-tailed Lizard). USA: FLORIDA: ST. LUCIE Co.: Port Saint Lucie, SW Biltmore Street & SW Molloy Street (27.28645°N, 80.36385°W, WGS84; elev. 5 m). 04 April 2010. Kenneth L. Krysko, Steve A. Johnson, Kyle E. Giddens, Kurt H. Gielow, Travis S. Lowke, William M. Moore, Eric Suarez, Cailey D. Thomas. Verified by Kevin M. Enge. Florida Museum of Natural History (UF 157453). New county record and extends the range ca. 44 km northwest of the closest known verified locality in Martin Co. (Hauge and Butterfield 2000. *Herpetol. Rev.* 31:53).

Submitted by **KENNETH L. KRYSKO** (e-mail: kenneyk@flmnh.ufl.edu), **KYLE E. GIDDENS**, **KURT H. GIELOW**, **TRAVIS S. LOWKE**, **WILLIAM M. MOORE**, **ERIC SUAREZ**, **CAILEY D. THOMAS**, Division of Herpetology, Florida Museum of Natural History, Dickinson Hall, University of Florida, Gainesville, Florida 32611, USA; and **STEVE A. JOHNSON**, Gulf Coast REC-Plant City Center and Department of Wildlife Ecology & Conservation, 1200 N. Park Road, Plant City, Florida 33563, USA.

OPHISAURUS ATTENUATUS ATTENUATUS (Western Slender Glass Lizard). USA: TEXAS: TYLER Co.: Big Thicket National Preserve, Turkey Creek Unit (30.58471°N, 94.33588°W; WGS 84). 28 November 2008. P. Crump, R. Rommel, and H. Pate. Verified by Carl Franklin. University of Texas Arlington UTADC 4015 – 4017. New county record (Dixon, J.R., 2000. *Amphibians and Reptiles of Texas*. 2nd ed. TAMU Press, College Station, Texas. 432 pp.).

Submitted by **PAUL CRUMP** (e-mail: pcrump@houstonzoo.org), **RACHEL ROMMEL** (e-mail: rrommel@houstonzoo.org), Department of Conservation and Science, Houston Zoo,

1513 North Macgregor Way, Houston, Texas 77030, USA; and **HAIGLER PATE**, National Park Service, Big Thicket National Preserve, 6044 FM 420, Kountze, Texas 77625, USA (e-mail: Haigler_Pate@nps.gov).

PLESTIODON LATICEPS (Broad-headed Skink). USA: ARKANSAS: OUACHITA Co.: ~23 km W of Camden (33.584058°N, 93.085246°W; WGS 84). AOR on Ouachita Co. Rd. 76, 1.5 km W Junction Co. Rd. 76/AR 57. 29 May 2010. K. Roberts and M. B. Connior. Verified by S. E. Trauth. Arkansas State University Museum of Zoology Herpetology Collection (photo voucher ASUMZ 31528). New county record (Trauth et al. 2004. *The Amphibians and Reptiles of Arkansas*. University of Arkansas Press, Fayetteville. 421 pp.). The species has also been collected from neighboring Columbia and Union counties in southern Arkansas.

Submitted by **MATTHEW B. CONNIOR**, Health and Natural Sciences, South Arkansas Community College, 300 S. West Avenue, El Dorado, Arkansas 71730, USA (e-mail: mconnior@southark.edu); **KORY ROBERTS**, Rogers High School, 2300 South Dixieland Road, Rogers, Arkansas 72758, USA; and **IDUN GUENTHER**, Fisheries and Wildlife Biology, Arkansas Tech University, Russellville, Arkansas 72801, USA.

SQUAMATA — SNAKES

AGKISTRODON CONTORTRIX MOKASEN (Northern Copperhead). USA: GEORGIA: WHITFIELD Co.: Heritage Point Park (34.7942972°N, 84.9445944°W; no datum available). 19 May 2010. Chris Manis and John Patrick. Verified by John Jensen. University of Tennessee at Chattanooga Museum of Natural History (UTC Digital Collection 4667TPW-GMU). New county record (Jensen et al. 2008. *Amphibians and Reptiles of Georgia*. University of Georgia Press, Athens. 570 pp.). Specimen was found in an open fairway of a canopied disc golf course.

Submitted by **CHRIS MANIS** and **THOMAS P. WILSON**, Department of Biological and Environmental Sciences, University of Tennessee at Chattanooga, 215 Holt Hall, Department 2653, 615 McCallie Avenue, Chattanooga, Tennessee 37403, USA; and **JOHN PATRICK**, Dalton Middle School, 1250 Cross Plains Trail, Dalton, Georgia 30721, USA.

AGKISTRODON PISCIVORUS LEUCOSTOMA (Western Cottonmouth). USA: TENNESSEE: LEWIS Co.: Buffalo River, 6.3 km downstream from Natchez Trace Parkway (35.4687°N, 87.5027°W; NAD 83). 10 July 2008. Todd Crabtree and Dwayne Estes. David H. Snyder Museum of Zoology, Austin Peay State University (APSU 18924, 18925, color photos). Verified by Nathan Parker. New county record (Scott and Redmond. 2008 [latest update: 14 July 2008]. *Atlas of Reptiles in Tennessee*. The Center for Field Biology, Austin Peay State University, Clarksville, Tennessee. Available from <http://apsu.edu/reptatlas/> [accessed 23 July 2008]). Range extension of ca. 150 river km up the Buffalo River drainage from the nearest downstream record in Humphreys Co. Adult on log overhanging water and juvenile on lower ledge of shale bluff along river; sky overcast with light rain, 1346 h.

Submitted by **TODD CRABTREE**, Division of Natural