

Creek. SCL 28 mm. First provincial record. An introduced species in the Philippines, previously recorded on Luzon, Mindoro, Panay, Cebu, Bohol, Leyte, and Mindanao (Diesmos et al. 2008. *Chelon. Conserv. Biol.* 7:157–177; Sy et al. 2004. *Turtle & Tortoise Newsl.* 7:17–18). Record extends distribution by ca. 309 km from Zamboanga City, the first record on Mindanao.

EMERSON Y. SY, Philippine Center for Terrestrial and Aquatic Research, 1198 Benavidez St., Unit 1202, Tondo, Manila, Philippines; e-mail: emersonsy@gmail.com.

PSEUDEMYX CONCINNA (River Cooter). USA: ALABAMA: CHOCTAW Co.: Two-track dirt road next to Okatuppa Creek (31.8306833°N, 88.2291138°W; WGS 84). 27 May 2008. Peter V. Lindeman. Verified by James C. Godwin. Florida Museum of Natural History (UF 153480 photographic voucher). New county record (Mount 1975. *The Reptiles and Amphibians of Alabama*. Auburn Univ. Agric. Exp. Sta., Auburn, Alabama. 347 pp.). Nesting adult female.

PETER V. LINDEMAN, Department of Biology and Health Services, 126 Cooper Hall, Edinboro University of Pennsylvania, Edinboro, Pennsylvania 16444, USA; e-mail plindeman@edinboro.edu.

PSEUDEMYX NELSONI (Florida Red-bellied Cooter). USA: FLORIDA: SUWANNEE Co.: Ichetucknee Springs State Park, Ichetucknee River near W bank 3 km NE of the Highway 27 bridge crossing (29.96922°N, 82.76069°W; WGS84). 22 February 2014. Jonathan D. Mays. Verified by Kenneth L. Krysko. Florida Museum of Natural History (UF 172555 photo voucher). New county record (Krysko et al. 2011. *Atlas of Amphibians and Reptiles in Florida*. Final report, Florida Fish and Wildlife Conservation Commission, Tallahassee. 524 pp.). I observed one basking adult while canoeing the spring run through a wide marshy area. A previous study reported *P. nelsoni* from the Ichetucknee (Chapin and Meylan 2010. *Herpetol. Conserv. Biol.* 6:51–60), but no museum vouchers were submitted for either Columbia or Suwannee counties, which are divided by the Ichetucknee River.

JONATHAN D. MAYS, Florida Fish and Wildlife Conservation Commission, 1105 S.W. Williston Road, Gainesville, Florida 32601, USA; e-mail: jonathan.mays@myfwc.com.

TRACHEMYS SCRIPTA ELEGANS (Red-eared Slider). USA: INDIANA: HAMILTON Co.: Conner Prairie (39.9820°N, 86.0309°W; WGS 84). 9 June 2013. Robert Brodman, Michael S. Finkler, James Horton, Megan Gramhofer, and Anastasia Marsh. Verified by Alan Resetar. Saint Joseph's College zoological collection (SJCZC R322). New county record (Minton 2001. *Amphibians & Reptiles of Indiana*, 2nd ed. Indiana Academy of Science, Indianapolis. 404 pp.). Extends range into a nine county gap 13 km N from nearest known locality in Marion County.

ROBERT BRODMAN (e-mail bobb@saintjoe.edu), **MEGAN GRAMHOFER** and **ANASTASIA MARSH**, Biology Department, Saint Joseph's College, Rensselaer, Indiana 47978, USA; **MICHAEL S. FINKLER**, School of Sciences, Indiana University Kokomo, PO Box 9003, Kokomo, Indiana 46904-9003, USA (e-mail: mfinkler@iuk.edu); **JAMES HORTON**, 10 Catalina Circle, Zionsville, Indiana 46077, USA (e-mail: stardali84@hotmail.com).

TRACHEMYS SCRIPTA SCRIPTA × ELEGANS (Pond Slider intergrade) USA: Pennsylvania: Berks Co.: Reading, Carsonia Park (40.337691°N, 75.872571°W; WGS84). 21 Sept 2013. Carl Maiese. Verified by Walter Meshaka. Kutztown University Museum of Zoology (KUMZ 1001). New county record (Hulse et al. 2001. *Amphibians and Reptiles of Pennsylvania*. Cornell University

Press, Ithaca, New York. 419 pp.). The online Pennsylvanian Amphibian and Reptile Survey (PARS, <http://www.paherpatlas.org> [accessed 8 Oct 2013]) noted *Trachemys scripta elegans* (online photo voucher only) and *Trachemys scripta* (no subspecific designation) observations, but *T. scripta scripta* and *T. scripta scripta* × *elegans* have not been confirmed from Berks Co. One *T. scripta* photographic entry from PARS suggested a possible *T. s. elegans* × *T. scripta* observation; however, our specimen is the first verified *T. s. scripta* × *T. s. elegans* intergrade documented in this county for which there is a preserved voucher. The male turtle was found in a pond in Carsonia Park; subsequent surveys uncovered an additional specimen and suggests that a breeding population of this introduced species has been established at this site.

CARL MAIESE (e-mail: cmaie796@live.kutztown.edu) and **MATTHEW STONE** (e-mail: stone@kutztown.edu), Department of Biology, Kutztown University, P.O. Box 730, Kutztown, Pennsylvania 19530, USA.

TRACHEMYS SCRIPTA SCRIPTA (Yellow-bellied Slider). USA: ALABAMA: BULLOCK Co.: basking in pond on the Wehle Tract, 8.4 km W of Comer, Alabama (32.03081°N 85.47115°W; WGS 84). 11 March 2014. C. Thawley and F. Graves. Verified by David Lauro. AUM AHAP-D 785–786 (digital photograph files). New county record (Mount 1975. *The Reptiles and Amphibians of Alabama*. Auburn Printing Co., Auburn, Alabama. 347 pp.). This record fills a distributional gap between the nearest vouchered localities approximately 10 km E in Barbour Co. and 29 km N in Macon Co.

CHRISTOPHER J. THAWLEY, Department of Biology, Pennsylvania State University, 208 Mueller Laboratory, University Park, Pennsylvania 16802, USA (e-mail: cthawley@gmail.com); **FERN GRAVES**, School of Forestry and Wildlife Sciences, Auburn University, Auburn, Alabama 36849, USA (e-mail: fbg0002@auburn.edu).

SQUAMATA — LIZARDS

AMPHISBAENA FULIGINOSA (Speckled Worm Lizard). BRAZIL: AMAPÁ: MUNICIPALITY OF LARANJAL DO JARI: Água Branca do Cajari (0.578314°S, 52.326769°W, 136 m elev.; datum WGS84) 10 July 2013. C. E. Costa-Campos, D. W. Silva-Silva, and K. J. Gaya Corrêa. Verified by A. P. Lima. Collection of Laboratório de Zoologia of the Universidade Federal do Amapá, Macapá, Amapá, Brazil (CDLABZOO 28, photo voucher). *Amphisbaena fuliginosa* was known previously from Panama, Peru, Ecuador, Colombia, Bolivia, Guyana, Suriname, Trinidad, and Brazil (Vanzolini 2002. *An. Acad. Bras. Ciênc.* 74[4]:609–648). In Brazil, the species was reported in the states of Pará (Ávila-Pires et al. 2010. *Bol. Mus. Para. Emílio Goeldi* 5[1]:13–112), Goiás (Colli et al. 2002. In Oliveira and Marquis [eds.], *The Cerrado of Brazil: Ecology and Natural History of a Neotropical Savanna*, pp. 223–241. Columbia University Press, New York), Minas Gerais (Silveira 2007. *Herpetol. Rev.* 38[4]:481; Nogueira-Costa et al. 2013. *Check List* 9[1]:167–168), Maranhão, Mato Grosso, Rondônia, Roraima, and Manaus (Vanzolini, *op. cit.*). First state record, extending the range 325 km NW from the municipality of Alenquer, Para state.

CARLOS E. COSTA-CAMPOS (e-mail: eduardocampos@unifap.br), **DANIEL WELTON SILVA-SILVA** (e-mail: dan.ws@hotmail.com), and **KELLY JULIANA GAYA CORRÊA** (e-mail: kelly_juliana@hotmail.com), Laboratório de Herpetologia, Departamento de Ciências Biológicas e da Saúde, Universidade Federal do Amapá, Campus Marco Zero, CEP 68903-419, Macapá, AP, Brazil.

AMPHISBAENA IGNATIANA. BRAZIL: PARÁÍBA: MUNICIPALITY OF JOÃO PESSOA: 7.1653°S, 34.8694°W. May 1984. Verified by S. da Silva.